# Proton number cumulants in heavy-ion collisions from hydrodynamics and the search for the QCD critical point

#### Volodymyr Vovchenko (INT Seattle / Berkeley Lab)

Seminar at the Department of High-Density Physics, BITP, Kyiv, Ukraine

#### June 22, 2022

VV, V. Koch, Phys. Rev. C 103, 044903 (2021)

VV, V. Koch, C. Shen, Phys. Rev. C 105, 014904 (2022)

VV, V. Koch, arXiv:2204.00137



## **QCD** phase structure



- Dilute hadron gas at low T &  $ho_{
  m B}$  due to confinement, quark-gluon plasma high T &  $ho_{
  m B}$
- Nuclear liquid-gas transition in cold and dense matter, lots of other phases conjectured

Is there a critical point and how to find it with heavy-ion collisions?

#### **Event-by-event fluctuations and statistical mechanics**



Cumulants measure chemical potential derivatives of the (QCD) equation of state

• (QCD) critical point – large correlation length, critical fluctuations of baryon number



M. Stephanov, PRL '09, '11 Energy scans at RHIC (STAR) and CERN-SPS (NA61/SHINE)

$$\kappa_2\sim\xi^2$$
,  $\kappa_3\sim\xi^{4.5}$ ,  $\kappa_4\sim\xi^7$ 

 $\xi \to \infty$ 

Looking for enhanced fluctuations and non-monotonicities

**Critical opalescence** 



#### **Example: Nuclear liquid-gas transition**



VV, Anchishkin, Gorenstein, Poberezhnyuk, PRC 92, 054901 (2015)

**Example: Lennard-Jones fluid** 

along the (super)critical isotherm of the liquid-gas transition

Microcanonical (const. EVN) ensemble with periodic boundary conditions

Variance of conserved particle number distribution inside coordinate space subvolume  $|z| < z^{max}$  as time average

$$ilde{\omega}^{\mathsf{coord}} = rac{1}{1-lpha} \, rac{\langle N^2 
angle - \langle N 
angle^2}{\langle N 
angle}$$







g.c.e.

1.00



N = 25000

N = 5000

N = 1000

N = 400

0.50

α

0.75

#### Measuring cumulants in heavy-ion collisions



Cumulants are extensive,  $\kappa_n \sim V$ , use ratios to cancel out the volume

$$\frac{\kappa_2}{\langle N \rangle}$$
,  $\frac{\kappa_3}{\kappa_2}$ ,  $\frac{\kappa_4}{\kappa_2}$ 

#### **Experimental measurements**

Beam energy scan in search for the critical point (STAR Coll.)



STAR Coll., Phys. Rev. Lett. 126, 092301 (2021); arXiv:2112.00240

Reduced errors (better statistics), more energies, to come soon from RHIC-BES-II program, STAR-FXT etc.

Can we learn more from the more accurate data available for  $\kappa_2$  and  $\kappa_3$ ?

Other measurements: LHC-ALICE, GSI-HADES & CERN-NA61/SHINE Collaborations

# **Theory vs experiment: Challenges for fluctuations**

#### Theory



 $\ensuremath{\mathbb{C}}$  Lattice QCD@BNL

- Coordinate space
- In contact with the heat bath
- Conserved charges
- Uniform
- Fixed volume

#### Experiment



STAR event display

- Momentum space
- Expanding in vacuum
- Non-conserved particle numbers
- Inhomogenous
- Fluctuating volume

#### Need dynamical description

# Dynamical approaches to the QCD critical point search

- 1. Dynamical model calculations of critical fluctuations
  - Fluctuating hydrodynamics
  - Equation of state with tunable critical point [P. Parotto et al, Phys. Rev. C 101, 034901 (2020)]
     Under development within the Beam Energy Scan Theory (BEST) Collaboration
     EEST [X. An et al., Nucl. Phys. A 1017, 122343 (2022)]
- 2. Molecular dynamics with a critical point

V. Kuznietsov et al., Phys. Rev. C 105, 044903 (2022)

- **3.** Deviations from precision calculations of non-critical fluctuations
  - Include essential non-critical contributions to (net-)proton number cumulants
  - Exact baryon conservation + hadronic interactions (hard core repulsion)
  - Based on realistic hydrodynamic simulations tuned to bulk data

[VV, C. Shen, V. Koch, Phys. Rev. C 105, 014904 (2022)]



## **Excluded volume effect**

Incorporate repulsive baryon (nucleon) hard core via excluded volume VV, M.I. Gorenstein, H. Stoecker, Phys. Rev. Lett. 118, 182301 (2017)

Amounts to a van der Waals correction for baryons in the HRG model

 $V \rightarrow V - bN$ 



 $\leftarrow 2r \rightarrow$ 





Figure from Ishii et al., PRL '07

• Net baryon kurtosis suppressed as in lattice QCD

$$\frac{\chi_4^B}{\chi_2^B} \simeq 1 - \frac{12b\phi_B(T)}{\Phi_B(T)} + O(b^2)$$

• Reproduces virial coefficients of baryon interaction from lattice QCD

Excluded volume from lattice QCD: b

$$b \approx 1 \text{ fm}^3$$



VV, A. Pasztor, S. Katz, Z. Fodor, H. Stoecker, Phys. Lett. B 755, 71 (2017) 10

# Hydrodynamic description within non-critical physics

- Collision geometry based 3D initial state
  - Constrained to net proton distributions [Shen, Alzhrani, Phys. Rev. C '20]
- Viscous hydrodynamics evolution MUSIC-3.0
  - Energy-momentum and baryon number conservation
  - Crossover equation of state based on lattice QCD [Monnai, Schenke, Shen, Phys. Rev. C '19]
- Cooper-Frye particlization at  $\epsilon_{sw} = 0.26 \text{ GeV}/\text{fm}^3$

$$\omega_p \frac{dN_j}{d^3 p} = \int_{\sigma(x)} d\sigma_\mu(x) p^\mu \frac{d_j \lambda_j^{\mathsf{ev}}(x)}{(2\pi)^3} \exp\left[\frac{\mu_j(x) - u^\mu(x)p_\mu}{T(x)}\right].$$

- Particlization respects QCD-based baryon number distribution
  - Incorporated via baryon excluded volume b = 1 fm<sup>3</sup>
     [VV, V. Koch, Phys. Rev. C 103, 044903 (2021)]
- Incorporates exact global baryon conservation via a method SAM-2.0



VV, V. Koch, C. Shen, Phys. Rev. C 105, 014904 (2022)





# **Calculating cumulants from hydrodynamics**

- Strategy:
  - 1. Calculate proton cumulants in the experimental acceptance in the grand-canonical limit
  - 2. Apply correction for the exact global baryon number conservation

First step:

- Sum contributions from each hypersurface element  $x_i$  at freeze-out
  - Cumulants of joint (anti)proton/(anti)baryon distribution

$$\kappa_{n,m}^{B^{\pm},p^{\pm},\text{gce}}(\Delta p_{\text{acc}}) = \sum_{i \in \sigma} \delta \kappa_{n,m}^{B^{\pm},p^{\pm},\text{gce}}(x_i;\Delta p_{\text{acc}}) \qquad \qquad p_{\text{acc}}(x_i;\Delta p_{\text{acc}}) = \frac{\int_{p \in \Delta p_{\text{acc}}} \frac{d^3p}{\omega_p} \delta \sigma_\mu(x_i) p^\mu f[u^\mu(x_i)p_\mu;T(x_i),\mu_j(x_i)]}{\int \frac{d^3p}{\omega_p} \delta \sigma_\mu(x_i) p^\mu f[u^\mu(x_i)p_\mu;T(x_i),\mu_j(x_i)]} \,.$$

- To compute each contribution
  - GCE susceptibilities  $\chi^{B^{\pm}}(x_i)$  define the distribution of the emitted (anti)baryons
  - Each baryon ends up in acceptance  $\Delta p_{acc}$  with binomial probability via the Cooper-Frye formula
  - Each baryon is a proton with probability  $q(x_i) = \langle N_p(x_i) \rangle / \langle N_B(x_i) \rangle$

## **Correcting for baryon number conservation with SAM-2.0**

$$P_1^{ ext{ce}}(B_1) \propto \sum_{B_1,B_2} P_1^{ ext{gce}}(B_1) P_2^{ ext{gce}}(B_2) imes rac{\delta_{B,B_1+B_2}}{\delta_{B,B_1+B_2}}$$

**SAM-1.0: uniform** thermal system and **coordinate** space

**SAM-2.0:** apply the correction for *arbitrary* distributions inside and outside the acceptance that are peaked at the mean

- Spatially inhomogeneous systems (e.g. RHIC)
- Momentum space
- Non-conserved quantities (e.g. proton number)
- Map "grand-canonical" cumulants inside and outside the acceptance to the "canonical" cumulants inside the acceptance

$$\kappa_{p,B}^{\text{in,ce}} = \mathsf{SAM}\left[\kappa_{p,B}^{\text{in,gce}}, \kappa_{p,B}^{\text{out,gce}}
ight]$$



VV, arXiv:2107.00163 (to appear in PRC)

# Net-particle fluctuations at the LHC (blast-wave)

- Net protons described within errors and consistent with either
  - global baryon conservation without  $B\overline{B}$  annihilations see e.g. ALICE Coll. arXiv:2206.03343
  - or local baryon conservation with  $B\overline{B}$  annihilations

O. Savchuk et al., Phys. Lett. B 827, 136983 (2022)



 Large effect from resonance decays for pions and kaons + exact conservation of electric charge/strangeness



VV, Koch, Phys. Rev. C 103, 044903 (2021) 14

# **RHIC-BES:** Net proton cumulant ratios (MUSIC)



- Data at  $\sqrt{s_{NN}} \ge 20$  GeV consistent with non-critical physics (baryon conservation and repulsion)
- Effect from baryon conservation is larger than from repulsion
- Excess of skewness in data at  $\sqrt{s_{NN}} < 20$  GeV hint of attractive interactions?

# **RHIC-BES:** Net proton cumulant ratios (MUSIC)



- Data at  $\sqrt{s_{NN}} \ge 20$  GeV consistent with non-critical physics (baryon conservation and repulsion)
- Effect from baryon conservation is larger than from repulsion
- Excess of skewness in data at  $\sqrt{s_{NN}} < 20$  GeV hint of attractive interactions?

# **Correlation Functions**

• Analyze genuine multi-particle correlations via factorial cumulants  $\hat{C}_n$ [Bzdak, Koch, Strodthoff, Phys. Rev. C '17]

$$\hat{C}_1 = \kappa_1, \qquad \hat{C}_3 = 2\kappa_1 - 3\kappa_2 + \kappa_3,$$

$$\hat{C}_2 = -\kappa_1 + \kappa_2, \qquad \hat{C}_4 = -6\kappa_1 + 11\kappa_2 - 6\kappa_3 + \kappa_4$$

$$\hat{C}_n^{\text{cons}} \propto \alpha^n, \qquad \hat{C}_n^{\text{EV}} \propto b^n$$
[Padels Keels Sheley EDIC [17]]

[Bzdak, Koch, Skokov, EPJC '17]

[VV et al, PLB '17]

- Three- and four-particle correlations are small without a CP
  - Multi-particle correlations expected near the critical point [Ling, Stephanov, PRC '15]
- Signals from the data at  $\sqrt{s_{NN}} \le 20$  GeV
  - Excess of two-proton correlations
  - Possibility of significant 4-proton correlations
  - Critical point?





• Changing  $y_{max}$  slope at  $\sqrt{s_{NN}} \le 14.5$  GeV?



- Changing  $y_{max}$  slope at  $\sqrt{s_{NN}} \le 14.5$  GeV?
- Volume fluctuations? [Skokov, Friman, Redlich, PRC '13]
  - $C_2/C_1 += C_1 * v_2$



- Changing  $y_{max}$  slope at  $\sqrt{s_{NN}} \le 14.5$  GeV?
- Volume fluctuations? [Skokov, Friman, Redlich, PRC '13]
  - $C_2/C_1 += C_1 * v_2$
  - Can improve low energies but spoil high energies?



- Changing  $y_{max}$  slope at  $\sqrt{s_{NN}} \le 14.5$  GeV?
- Volume fluctuations? [Skokov, Friman, Redlich, PRC '13]
  - $C_2/C_1 += C_1 * v_2$
  - Can improve low energies but spoil high energies?
- Exact electric charge conservation?
  - Worsens the agreement at  $\sqrt{s_{NN}} \leq 14.5\,,$  higher energies virtually unaffected



- Changing  $y_{max}$  slope at  $\sqrt{s_{NN}} \le 14.5$  GeV? ٠
- Volume fluctuations? [Skokov, Friman, Redlich, PRC '13] ٠
  - $C_2/C_1 + = C_1 * v_2$ •
  - Can improve low energies but spoil high energies? •
- Exact electric charge conservation? •
  - Worsens the agreement at  $\sqrt{s_{NN}} \le 14.5$ , higher energies ٠ virtually unaffected

⊢ <sub>15</sub>

das

μ (MeV)

- **Attractive interactions?** 
  - Could work if baryon repulsion turns • into attraction in the high- $\mu_B$  regime (MeV) 20
  - **Critical point?** •



# Lower energies $\sqrt{s_{NN}} \le 7.7$ GeV



 Fill the gap with ongoing/future data from STAR-FXT (e.g. arXiv:2112.00240), future experiments like CBM-FAIR

# Thermodynamic analysis of HADES data

- Single freeze-out scenario: Emission from Siemens-Rasmussen hypersurface with Hubblelike flow
  - $\rightarrow$  Pion and proton spectra o.k. [S. Harabasz et al., PRC 102, 054903 (2020)]
- Uniform  $T \approx 70$  MeV,  $\mu_B \approx 875$  MeV across the fireball [A. Motornenko et al., PLB 822, 136703 (2021)]
- Fluctuations:
  - Same as before but incorporate additional binomial filtering to account for protons bound in light nuclei
  - Uniform fireball  $\rightarrow$  Final proton cumulants are linear combinations of baryon susceptibilities  $\chi^B_n$  at freezeout





- Fit baryon susceptibilities to data within a fireball model (Siemens-Rasmussen\*)
- In the grand-canonical limit (no baryon conservation, small  $y_{cut}$ ) the data are described well with

$$\frac{\chi_2^B}{\chi_1^B} = 9.35 \pm 0.40, \qquad \frac{\chi_3^B}{\chi_2^B} = -39.6 \pm 7.2, \qquad \frac{\chi_4^B}{\chi_2^B} = 1130 \pm 488 \qquad \text{i.e.} \qquad \chi_4^B \gg -\chi_3^B \gg \chi_2^B \gg \chi_1^B$$

- Could be indicative of a *critical point* near the HADES freeze-out at  $T \sim 70$  MeV,  $\mu_B \sim 875$  MeV
- However, the results for  $y_{cut} > 0.2$  are challenging to describe with baryon conservation included



## Summary: What we learned so far from fluctuations



- Data at high energies ( $\sqrt{s_{NN}} \ge 20$  GeV) consistent with "non-critical" physics
  - Disfavors QCD critical point at  $\mu_B/T < 2-3$ , consistent with what we know from lattice QCD
- Interesting indications for (multi)-proton correlations at  $\sqrt{s_{NN}} \le 7.7$  GeV

#### Thanks for your attention!